



United States
Department of
Agriculture

NRCS

Natural
Resources
Conservation
Service

A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for **Bladen County, North Carolina**

Kelly Farm and Hunting Land



October 29, 2025

Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<https://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	8
Soil Map.....	9
Legend.....	10
Map Unit Legend.....	11
Map Unit Descriptions.....	11
Bladen County, North Carolina.....	13
Ce—Centenary sand.....	13
Cr—Croatan muck, ponded, 0 to 2 percent slopes.....	14
KeA—Kenansville sand, 0 to 3 percent slopes.....	15
LaB—Lakeland sand, 1 to 7 percent slopes.....	16
LeA—Leon sand, 0 to 3 percent slopes.....	17
Ly—Lynn Haven and Torhunta soils.....	18
Pp—Paxville sandy loam.....	20
Wh—Wasda muck.....	22
Soil Information for All Uses	24
Suitabilities and Limitations for Use.....	24
Vegetative Productivity.....	24
Forest Productivity (Tree Site Index): longleaf pine (USDA 1929 (580)).....	24
Forest Productivity (Tree Site Index): loblolly pine (Coile, Schumacher 1953 (690)).....	27
Yields of Non-Irrigated Crops (Component): Soybeans (Bu).....	30
Yields of Non-Irrigated Crops (Component): Corn (Bu).....	34
References	39

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and


Custom Soil Resource Report

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map



Custom Soil Resource Report

MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

 Soil Map Unit Polygons

 Soil Map Unit Lines

 Soil Map Unit Points

Special Point Features

 Blowout

 Borrow Pit

 Clay Spot

 Closed Depression

 Gravel Pit

 Gravelly Spot

 Landfill

 Lava Flow

 Marsh or swamp

 Mine or Quarry

 Miscellaneous Water

 Perennial Water

 Rock Outcrop

 Saline Spot

 Sandy Spot

 Severely Eroded Spot

 Sinkhole

 Slide or Slip

 Sodic Spot

 Spoil Area

 Stony Spot

 Very Stony Spot

 Wet Spot

 Other

 Special Line Features

Water Features

 Streams and Canals

Transportation

 Rails

 Interstate Highways

 US Routes

 Major Roads

 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
Web Soil Survey URL:
Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Bladen County, North Carolina
Survey Area Data: Version 29, Sep 2, 2025

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Apr 11, 2022—May 15, 2022

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
Ce	Centenary sand	2.1	1.5%
Cr	Croatan muck, ponded, 0 to 2 percent slopes	12.4	8.8%
KeA	Kenansville sand, 0 to 3 percent slopes	3.9	2.8%
LaB	Lakeland sand, 1 to 7 percent slopes	24.8	17.5%
LeA	Leon sand, 0 to 3 percent slopes	14.1	10.0%
Ly	Lynn Haven and Torhunta soils	15.0	10.6%
Pp	Paxville sandy loam	4.5	3.2%
Wh	Wasda muck	64.4	45.6%
Totals for Area of Interest		141.2	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not

mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Bladen County, North Carolina

Ce—Centenary sand

Map Unit Setting

National map unit symbol: 3w4x
Elevation: 80 to 330 feet
Mean annual precipitation: 38 to 55 inches
Mean annual air temperature: 59 to 70 degrees F
Frost-free period: 210 to 265 days
Farmland classification: Not prime farmland

Map Unit Composition

Centenary and similar soils: 75 percent
Minor components: 5 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Centenary

Setting

Landform: Flats on marine terraces, ridges on marine terraces
Down-slope shape: Convex
Across-slope shape: Concave
Parent material: Sandy fluviomarine deposits and/or eolian sands

Typical profile

A - 0 to 9 inches: sand
E - 9 to 58 inches: sand
Bh - 58 to 80 inches: sand

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Very low
Capacity of the most limiting layer to transmit water (Ksat): High to very high (1.98 to 19.98 in/hr)
Depth to water table: About 36 to 60 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 4.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3s
Hydrologic Soil Group: A
Ecological site: F153AY050NC - Moist Spodosol Rises and Flats
Hydric soil rating: No

Minor Components

Leon

Percent of map unit: 5 percent
Landform: Flats on marine terraces
Down-slope shape: Linear
Across-slope shape: Concave

Custom Soil Resource Report

Ecological site: F153AY070NC - Wet Spodosol Flats and Depressions
Hydric soil rating: Yes

Cr—Croatan muck, ponded, 0 to 2 percent slopes

Map Unit Setting

National map unit symbol: 2tn9n
Elevation: 20 to 100 feet
Mean annual precipitation: 40 to 55 inches
Mean annual air temperature: 59 to 70 degrees F
Frost-free period: 200 to 280 days
Farmland classification: Not prime farmland

Map Unit Composition

Croatan and similar soils: 87 percent
Minor components: 13 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Croatan

Setting

Landform: Bays (geom.), carolina bays, flats, pocosins
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Herbaceous organic material over loamy fluviomarine deposits

Typical profile

Oa - 0 to 28 inches: muck
Ag - 28 to 33 inches: mucky sandy loam
Cg1 - 33 to 60 inches: sandy clay loam
Cg2 - 60 to 80 inches: loamy sand

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Very poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Very low to high (0.00 to 1.98 in/hr)
Depth to water table: About 0 to 6 inches
Frequency of flooding: None
Frequency of ponding: Frequent
Available water supply, 0 to 60 inches: Very high (about 16.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 7w
Hydrologic Soil Group: B/D
Ecological site: F153AY100NC - Flooded Organic Soil Flood Plains and Terraces
Hydric soil rating: Yes

Minor Components

Torhunta

Percent of map unit: 8 percent

Landform: Carolina bays on marine terraces, depressions on stream terraces, flats

Landform position (three-dimensional): Tread, dip, tal

Down-slope shape: Concave, linear

Across-slope shape: Concave, linear

Ecological site: F153AY090NC - Flooded Mineral Soil Flood Plains and Terraces

Hydric soil rating: Yes

Pantego

Percent of map unit: 5 percent

Landform: Depressions, broad interstream divides on marine terraces, flats on marine terraces

Landform position (three-dimensional): Tread, dip, tal

Down-slope shape: Concave, linear

Across-slope shape: Concave, linear

Ecological site: F153AY060NC - Wet Loamy Flats and Depressions

Hydric soil rating: Yes

KeA—Kenansville sand, 0 to 3 percent slopes

Map Unit Setting

National map unit symbol: 3w5n

Elevation: 80 to 330 feet

Mean annual precipitation: 38 to 55 inches

Mean annual air temperature: 59 to 70 degrees F

Frost-free period: 210 to 265 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Kenansville and similar soils: 95 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Kenansville

Setting

Landform: Stream terraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Loamy alluvium over sandy alluvium

Typical profile

Ap - 0 to 8 inches: loamy sand

E - 8 to 24 inches: loamy sand

Bt - 24 to 36 inches: sandy loam

BC - 36 to 42 inches: loamy sand

C - 42 to 84 inches: sand

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Very low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 5.95 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 4.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 2s
Hydrologic Soil Group: A
Ecological site: F153AY030NC - Dry Loamy Rises and Flats, F133AY220NC -
Stream Terraces, Moist
Hydric soil rating: No

LaB—Lakeland sand, 1 to 7 percent slopes

Map Unit Setting

National map unit symbol: 3w5q
Elevation: 80 to 330 feet
Mean annual precipitation: 38 to 55 inches
Mean annual air temperature: 59 to 70 degrees F
Frost-free period: 210 to 265 days
Farmland classification: Not prime farmland

Map Unit Composition

Lakeland and similar soils: 80 percent
Minor components: 5 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lakeland

Setting

Landform: Ridges on marine terraces
Landform position (two-dimensional): Shoulder, summit
Landform position (three-dimensional): Crest
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Sandy marine deposits and/or eolian sands

Typical profile

A - 0 to 6 inches: sand
C1 - 6 to 48 inches: sand
C2 - 48 to 80 inches: sand

Properties and qualities

Slope: 0 to 6 percent

Custom Soil Resource Report

Depth to restrictive feature: More than 80 inches
Drainage class: Excessively drained
Runoff class: Very low
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 4.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4s
Hydrologic Soil Group: A
Ecological site: F133AY720NC - Sandy Upland, Dry Summits, F153AY010NC - Dry Sands, F133AY240NC - Sandy Stream Terraces, Dry
Hydric soil rating: No

Minor Components

Leon

Percent of map unit: 5 percent
Landform: Flats on marine terraces
Down-slope shape: Linear
Across-slope shape: Concave
Ecological site: F153AY070NC - Wet Spodosol Flats and Depressions, F133AY340NC - Upland Depressions, Flats, and Carolina Bay Rims, Sandy
Hydric soil rating: Yes

LeA—Leon sand, 0 to 3 percent slopes

Map Unit Setting

National map unit symbol: 3w5r
Elevation: 80 to 330 feet
Mean annual precipitation: 38 to 55 inches
Mean annual air temperature: 59 to 70 degrees F
Frost-free period: 210 to 265 days
Farmland classification: Farmland of unique importance

Map Unit Composition

Leon and similar soils: 80 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Leon

Setting

Landform: Flats on marine terraces
Down-slope shape: Linear
Across-slope shape: Concave
Parent material: Sandy fluviomarine deposits and/or eolian sands

Typical profile

A - 0 to 3 inches: sand
E - 3 to 15 inches: sand
Bh - 15 to 30 inches: fine sand
BE - 30 to 33 inches: fine sand
E' - 33 to 66 inches: fine sand
B'h - 66 to 80 inches: fine sand

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Poorly drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.20 to 1.98 in/hr)
Depth to water table: About 0 to 12 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 3.8 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: A/D
Ecological site: F153AY070NC - Wet Spodosol Flats and Depressions
Hydric soil rating: Yes

Ly—Lynn Haven and Torhunta soils

Map Unit Setting

National map unit symbol: 3w5t
Elevation: 80 to 330 feet
Mean annual precipitation: 38 to 55 inches
Mean annual air temperature: 59 to 70 degrees F
Frost-free period: 210 to 265 days
Farmland classification: Prime farmland if drained

Map Unit Composition

Lynn haven, undrained, and similar soils: 41 percent
Torhunta, undrained, and similar soils: 39 percent
Torhunta, drained, and similar soils: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lynn Haven, Undrained

Setting

Landform: Flats on marine terraces
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Sandy fluviomarine deposits and/or eolian sands

Typical profile

A - 0 to 9 inches: sand
E - 9 to 12 inches: sand
Bh - 12 to 80 inches: sand

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Poorly drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 5.95 in/hr)
Depth to water table: About 0 to 6 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water supply, 0 to 60 inches: Very high (about 12.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: A/D
Ecological site: F153AY070NC - Wet Spodosol Flats and Depressions,
F133AY340NC - Upland Depressions, Flats, and Carolina Bay Rims, Sandy
Hydric soil rating: Yes

Description of Torhunta, Undrained

Setting

Landform: Carolina bays on marine terraces, flats on marine terraces, depressions on stream terraces
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Sandy and loamy alluvium and/or fluviomarine deposits

Typical profile

A - 0 to 15 inches: fine sandy loam
Bg - 15 to 40 inches: fine sandy loam
Cg - 40 to 80 inches: loamy sand

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Very poorly drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 5.95 in/hr)
Depth to water table: About 0 to 12 inches
Frequency of flooding: Rare
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 5.8 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 6w
Hydrologic Soil Group: A/D

Custom Soil Resource Report

Ecological site: F153AY090NC - Flooded Mineral Soil Flood Plains and Terraces,
F133AY320NC - Upland Depressions, Flats, and Carolina Bays
Hydric soil rating: Yes

Description of Torhunta, Drained

Setting

Landform: Depressions on stream terraces, carolina bays on marine terraces, flats on marine terraces
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Sandy and loamy alluvium and/or fluviomarine deposits

Typical profile

Ap - 0 to 15 inches: fine sandy loam
Bg - 15 to 40 inches: fine sandy loam
Cg - 40 to 80 inches: loamy sand

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Very poorly drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 5.95 in/hr)
Depth to water table: About 0 to 12 inches
Frequency of flooding: Rare
Frequency of ponding: None
Available water supply, 0 to 60 inches: Low (about 5.8 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3w
Hydrologic Soil Group: A/D
Ecological site: F133AY320NC - Upland Depressions, Flats, and Carolina Bays,
F153AY090NC - Flooded Mineral Soil Flood Plains and Terraces
Hydric soil rating: Yes

Pp—Paxville sandy loam

Map Unit Setting

National map unit symbol: 3w63
Elevation: 20 to 160 feet
Mean annual precipitation: 40 to 55 inches
Mean annual air temperature: 59 to 70 degrees F
Frost-free period: 200 to 280 days
Farmland classification: Prime farmland if drained

Map Unit Composition

Paxville, drained, and similar soils: 80 percent

Custom Soil Resource Report

Paxville, undrained, and similar soils: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Paxville, Drained

Setting

Landform: Broad interstream divides on marine terraces, carolina bays on marine terraces, flats on marine terraces

Landform position (two-dimensional): Summit

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Loamy marine deposits

Typical profile

Ap - 0 to 14 inches: sandy loam

Btg - 14 to 42 inches: sandy clay loam

BCg - 42 to 47 inches: sandy loam

Cg - 47 to 80 inches: sand

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 0 inches

Frequency of flooding: Rare

Frequency of ponding: None

Available water supply, 0 to 60 inches: Moderate (about 7.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3w

Hydrologic Soil Group: A/D

Ecological site: F153AY060NC - Wet Loamy Flats and Depressions

Hydric soil rating: Yes

Description of Paxville, Undrained

Setting

Landform: Broad interstream divides on marine terraces, carolina bays on marine terraces, flats on marine terraces

Landform position (two-dimensional): Summit

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Loamy marine deposits

Typical profile

A - 0 to 14 inches: sandy loam

Btg - 14 to 42 inches: sandy clay loam

BCg - 42 to 47 inches: sandy loam

Cg - 47 to 80 inches: sand

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Custom Soil Resource Report

Runoff class: Negligible

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: About 0 inches

Frequency of flooding: Rare

Frequency of ponding: Frequent

Available water supply, 0 to 60 inches: Moderate (about 7.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6w

Hydrologic Soil Group: A/D

Ecological site: F153AY060NC - Wet Loamy Flats and Depressions

Hydric soil rating: Yes

Wh—Wasda muck

Map Unit Setting

National map unit symbol: 3w6k

Elevation: 20 to 160 feet

Mean annual precipitation: 40 to 55 inches

Mean annual air temperature: 59 to 70 degrees F

Frost-free period: 200 to 280 days

Farmland classification: Prime farmland if drained

Map Unit Composition

Wasda, drained, and similar soils: 80 percent

Wasda, undrained, and similar soils: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Wasda, Drained

Setting

Landform: Depressions on marine terraces, flats on marine terraces

Down-slope shape: Convex

Across-slope shape: Concave

Parent material: Herbaceous organic material and/or woody organic material over
loamy marine deposits

Typical profile

Oap - 0 to 14 inches: muck

A - 14 to 20 inches: loam

Bg - 20 to 43 inches: clay loam

Cg1 - 43 to 60 inches: sandy loam

Cg2 - 60 to 80 inches: sand

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Runoff class: Very high

Custom Soil Resource Report

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)

Depth to water table: About 0 to 12 inches

Frequency of flooding: Rare

Frequency of ponding: None

Available water supply, 0 to 60 inches: High (about 10.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3w

Hydrologic Soil Group: C/D

Ecological site: F153AY060NC - Wet Loamy Flats and Depressions,

F153BY060NC - Wet Loamy Flats and Depressions

Hydric soil rating: Yes

Description of Wasda, Undrained

Setting

Landform: Depressions on marine terraces, flats on marine terraces

Down-slope shape: Convex

Across-slope shape: Concave

Parent material: Herbaceous organic material and/or woody organic material over loamy marine deposits

Typical profile

Oa - 0 to 14 inches: muck

A - 14 to 20 inches: loam

Bg - 20 to 43 inches: clay loam

Cg1 - 43 to 60 inches: sandy loam

Cg2 - 60 to 80 inches: sand

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)

Depth to water table: About 0 to 12 inches

Frequency of flooding: Rare

Frequency of ponding: None

Available water supply, 0 to 60 inches: High (about 10.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6w

Hydrologic Soil Group: C/D

Ecological site: F153AY060NC - Wet Loamy Flats and Depressions,

F153BY060NC - Wet Loamy Flats and Depressions

Hydric soil rating: Yes

Soil Information for All Uses

Suitabilities and Limitations for Use

The Suitabilities and Limitations for Use section includes various soil interpretations displayed as thematic maps with a summary table for the soil map units in the selected area of interest. A single value or rating for each map unit is generated by aggregating the interpretive ratings of individual map unit components. This aggregation process is defined for each interpretation.

Vegetative Productivity


Vegetative productivity includes estimates of potential vegetative production for a variety of land uses, including cropland, forestland, hayland, pastureland, horticulture and rangeland. In the underlying database, some states maintain crop yield data by individual map unit component. Other states maintain the data at the map unit level. Attributes are included for both, although only one or the other is likely to contain data for any given geographic area. For other land uses, productivity data is shown only at the map unit component level. Examples include potential crop yields under irrigated and nonirrigated conditions, forest productivity, forest site index, and total rangeland production under of normal, favorable and unfavorable conditions.

Forest Productivity (Tree Site Index): longleaf pine (USDA 1929 (580))

The "site index" is the average height, in feet, that dominant and codominant trees of a given species attain in a specified number of years. The site index applies to fully stocked, even-aged, unmanaged stands.

This attribute is actually recorded as three separate values in the database. A low value and a high value indicate the range of this attribute for the soil component. A "representative" value indicates the expected value of this attribute for the component. For this attribute, only the representative value is used.

Custom Soil Resource Report
Map—Forest Productivity (Tree Site Index): longleaf pine (USDA 1929 (580))



MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

Soil Rating Polygons

 ≤ 60
 > 60 and ≤ 65
 > 65 and ≤ 70
 > 70 and ≤ 72
 Not rated or not available

Soil Rating Lines

 ≤ 60
 > 60 and ≤ 65
 > 65 and ≤ 70
 > 70 and ≤ 72
 Not rated or not available

Soil Rating Points

 ≤ 60
 > 60 and ≤ 65
 > 65 and ≤ 70
 > 70 and ≤ 72
 Not rated or not available

Water Features

 Streams and Canals

Transportation

 Rails
 Interstate Highways

 US Routes
 Major Roads
 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Bladen County, North Carolina
 Survey Area Data: Version 29, Sep 2, 2025

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Apr 11, 2022—May 15, 2022

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Table—Forest Productivity (Tree Site Index): longleaf pine (USDA 1929 (580))

Map unit symbol	Map unit name	Rating (feet)	Acres in AOI	Percent of AOI
Ce	Centenary sand	72	2.1	1.5%
Cr	Croatan muck, ponded, 0 to 2 percent slopes		12.4	8.8%
KeA	Kenansville sand, 0 to 3 percent slopes	65	3.9	2.8%
LaB	Lakeland sand, 1 to 7 percent slopes	60	24.8	17.5%
LeA	Leon sand, 0 to 3 percent slopes	70	14.1	10.0%
Ly	Lynn Haven and Torhunta soils	70	15.0	10.6%
Pp	Paxville sandy loam		4.5	3.2%
Wh	Wasda muck		64.4	45.6%
Totals for Area of Interest			141.2	100.0%

Rating Options—Forest Productivity (Tree Site Index): longleaf pine (USDA 1929 (580))

Units of Measure: feet

Tree: longleaf pine

Site Index Base: USDA 1929 (580)

Aggregation Method: Dominant Component

Component Percent Cutoff: None Specified

Tie-break Rule: Higher

Interpret Nulls as Zero: No


Forest Productivity (Tree Site Index): loblolly pine (Coile, Schumacher 1953 (690))

The "site index" is the average height, in feet, that dominant and codominant trees of a given species attain in a specified number of years. The site index applies to fully stocked, even-aged, unmanaged stands.

This attribute is actually recorded as three separate values in the database. A low value and a high value indicate the range of this attribute for the soil component. A "representative" value indicates the expected value of this attribute for the component. For this attribute, only the representative value is used.

Custom Soil Resource Report

Map—Forest Productivity (Tree Site Index): loblolly pine (Coile, Schumacher 1953 (690))



MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

Soil Rating Polygons

 ≤ 75
 > 75 and ≤ 80
 > 80 and ≤ 85
 > 85 and ≤ 96
 Not rated or not available

Soil Rating Lines

 ≤ 75
 > 75 and ≤ 80
 > 80 and ≤ 85
 > 85 and ≤ 96
 Not rated or not available

Soil Rating Points

 ≤ 75
 > 75 and ≤ 80
 > 80 and ≤ 85
 > 85 and ≤ 96
 Not rated or not available

Water Features

 Streams and Canals

Transportation

 Rails
 Interstate Highways

 US Routes
 Major Roads
 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Bladen County, North Carolina
 Survey Area Data: Version 29, Sep 2, 2025

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Apr 11, 2022—May 15, 2022

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Table—Forest Productivity (Tree Site Index): loblolly pine (Coile, Schumacher 1953 (690))

Map unit symbol	Map unit name	Rating (feet)	Acres in AOI	Percent of AOI
Ce	Centenary sand	85	2.1	1.5%
Cr	Croatan muck, ponded, 0 to 2 percent slopes		12.4	8.8%
KeA	Kenansville sand, 0 to 3 percent slopes	80	3.9	2.8%
LaB	Lakeland sand, 1 to 7 percent slopes	75	24.8	17.5%
LeA	Leon sand, 0 to 3 percent slopes	75	14.1	10.0%
Ly	Lynn Haven and Torhunta soils	80	15.0	10.6%
Pp	Paxville sandy loam	96	4.5	3.2%
Wh	Wasda muck		64.4	45.6%
Totals for Area of Interest			141.2	100.0%

Rating Options—Forest Productivity (Tree Site Index): loblolly pine (Coile, Schumacher 1953 (690))*Units of Measure:* feet*Tree:* loblolly pine*Site Index Base:* Coile, Schumacher 1953 (690)*Aggregation Method:* Dominant Component*Component Percent Cutoff:* None Specified*Tie-break Rule:* Higher*Interpret Nulls as Zero:* No**Yields of Non-Irrigated Crops (Component): Soybeans (Bu)**

These are the estimated average yields per acre that can be expected of selected nonirrigated crops under a high level of management. In any given year, yields may be higher or lower than those indicated because of variations in rainfall and other climatic factors.

In the database, some states maintain crop yield data by individual map unit component and others maintain the data at the map unit level. Attributes are included in this application for both, although only one or the other is likely to

Custom Soil Resource Report

contain data for any given geographic area. This attribute uses data maintained at the map unit component level.


The yields are actually recorded as three separate values in the database. A low value and a high value indicate the range for the soil component. A "representative" value indicates the expected value for the component. For these yields, only the representative value is used.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby areas and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

The estimated yields reflect the productive capacity of each soil for the selected crop. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Custom Soil Resource Report
 Map—Yields of Non-Irrigated Crops (Component): Soybeans (Bu)



MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

Soil Rating Polygons

 ≤ 14.73
 > 14.73 and ≤ 26.00
 > 26.00 and ≤ 30.00
 > 30.00 and ≤ 35.56
 > 35.56 and ≤ 40.66
 Not rated or not available

Soil Rating Lines

 ≤ 14.73
 > 14.73 and ≤ 26.00
 > 26.00 and ≤ 30.00
 > 30.00 and ≤ 35.56
 > 35.56 and ≤ 40.66
 Not rated or not available

Soil Rating Points

 ≤ 14.73
 > 14.73 and ≤ 26.00
 > 26.00 and ≤ 30.00
 > 30.00 and ≤ 35.56
 > 35.56 and ≤ 40.66
 Not rated or not available

Water Features

 Streams and Canals

Transportation

 Rails
 Interstate Highways
 US Routes
 Major Roads
 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Bladen County, North Carolina
 Survey Area Data: Version 29, Sep 2, 2025

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Apr 11, 2022—May 15, 2022

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Table—Yields of Non-Irrigated Crops (Component): Soybeans (Bu)

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
Ce	Centenary sand	23.44	2.1	1.5%
Cr	Croatan muck, ponded, 0 to 2 percent slopes	40.66	12.4	8.8%
KeA	Kenansville sand, 0 to 3 percent slopes	30.00	3.9	2.8%
LaB	Lakeland sand, 1 to 7 percent slopes	22.24	24.8	17.5%
LeA	Leon sand, 0 to 3 percent slopes	26.00	14.1	10.0%
Ly	Lynn Haven and Torhunta soils	14.73	15.0	10.6%
Pp	Paxville sandy loam	35.56	4.5	3.2%
Wh	Wasda muck	40.00	64.4	45.6%
Totals for Area of Interest			141.2	100.0%

Rating Options—Yields of Non-Irrigated Crops (Component): Soybeans (Bu)*Crop:* Soybeans*Yield Units:* Bu*Aggregation Method:* Weighted Average*Component Percent Cutoff:* None Specified*Tie-break Rule:* Higher*Interpret Nulls as Zero:* Yes**Yields of Non-Irrigated Crops (Component): Corn (Bu)**

These are the estimated average yields per acre that can be expected of selected nonirrigated crops under a high level of management. In any given year, yields may be higher or lower than those indicated because of variations in rainfall and other climatic factors.

In the database, some states maintain crop yield data by individual map unit component and others maintain the data at the map unit level. Attributes are included in this application for both, although only one or the other is likely to contain data for any given geographic area. This attribute uses data maintained at the map unit component level.

Custom Soil Resource Report

The yields are actually recorded as three separate values in the database. A low value and a high value indicate the range for the soil component. A "representative" value indicates the expected value for the component. For these yields, only the representative value is used.


The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby areas and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

The estimated yields reflect the productive capacity of each soil for the selected crop. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Custom Soil Resource Report

Map—Yields of Non-Irrigated Crops (Component): Corn (Bu)



MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)

Soils

Soil Rating Polygons

 ≤ 39.67
 > 39.67 and ≤ 60.94
 > 60.94 and ≤ 80.00
 > 80.00 and ≤ 102.22
 > 102.22 and ≤ 124.67
 Not rated or not available

Soil Rating Lines

 ≤ 39.67
 > 39.67 and ≤ 60.94
 > 60.94 and ≤ 80.00
 > 80.00 and ≤ 102.22
 > 102.22 and ≤ 124.67
 Not rated or not available

Soil Rating Points

 ≤ 39.67
 > 39.67 and ≤ 60.94
 > 60.94 and ≤ 80.00
 > 80.00 and ≤ 102.22
 > 102.22 and ≤ 124.67
 Not rated or not available

Water Features

 Streams and Canals

Transportation

 Rails
 Interstate Highways
 US Routes
 Major Roads
 Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Bladen County, North Carolina
 Survey Area Data: Version 29, Sep 2, 2025

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Apr 11, 2022—May 15, 2022

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Table—Yields of Non-Irrigated Crops (Component): Corn (Bu)

Map unit symbol	Map unit name	Rating	Acres in AOI	Percent of AOI
Ce	Centenary sand	60.94	2.1	1.5%
Cr	Croatan muck, ponded, 0 to 2 percent slopes	124.67	12.4	8.8%
KeA	Kenansville sand, 0 to 3 percent slopes	80.00	3.9	2.8%
LaB	Lakeland sand, 1 to 7 percent slopes	54.94	24.8	17.5%
LeA	Leon sand, 0 to 3 percent slopes	70.00	14.1	10.0%
Ly	Lynn Haven and Torhunta soils	39.67	15.0	10.6%
Pp	Paxville sandy loam	102.22	4.5	3.2%
Wh	Wasda muck	120.00	64.4	45.6%
Totals for Area of Interest			141.2	100.0%

**Rating Options—Yields of Non-Irrigated Crops (Component):
Corn (Bu)**

Crop: Corn

Yield Units: Bu

Aggregation Method: Weighted Average

Component Percent Cutoff: None Specified

Tie-break Rule: Higher

Interpret Nulls as Zero: Yes

References

- American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.
- American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.
- Federal Register. July 13, 1994. Changes in hydric soils of the United States.
- Federal Register. September 18, 2002. Hydric soils of the United States.
- Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.
- National Research Council. 1995. Wetlands: Characteristics and boundaries.
- Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_054262
- Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577
- Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580
- Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.
- United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.
- United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053374
- United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. <http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084>

Custom Soil Resource Report

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf